

SEQUENCE LISTING

<110> Pietras, Kristian
Ostman, Arne
Heldin, Carl-Henrik
Rubin, Kristofer

<120> METHOD FOR TREATMENT OF TUMORS USING NUCLEIC ACID
LIGANDS TO PDGF

<130> NEX90

<140> 09/859,724
<141> 2001-05-17

<150> 60/205,006
<151> 2000-05-17

<150> 08/479,725
<151> 1995-06-07

<150> 08/479,783
<151> 1995-06-07

<150> 08/618,693
<151> 1996-03-20

<150> 08/991,743
<151> 1997-12-16

<160> 2

<170> PatentIn Ver. 2.0

<210> 1
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Nucleic Acid Ligand

<220>
<221> modified_base
<222> (6)..(30)
<223> U at positions 6, 20 and 30 is
2'-fluoro-2'-deoxyuridine

<220>
 <221> modified_base
 <222> (8)..(29)
 <223> C at positions 8, 21, 28, and 29 is
 2'-fluoro-2'-deoxycytidine

 <220>
 <221> modified_base
 <222> (9)..(31)
 <223> G at positions 9, 15, 17, and 31 is
 2'-O-Methyl-2'-deoxyguanosine

 <220>
 <221> modified_base
 <222> (22)
 <223> A at position 22 is 2'-O-Methyl-2'-deoxyadenosine

 <220>
 <221> modified_base
 <222> (1)..(30)
 <223> The residues at positions 9 and 10 are connected
 by a hexaethylene glycol phosphoramidite linker.
 The residues at positions 21 and 22 are
 connected by a hexaethylene glycol phosphoramidite
 linker.

 <220>
 <221> modified_base
 <222> (32)
 <223> Nucleotide 32 is an inverted orientation T
 (3'-3'-linked)

<400> 1
 caggcuacgc gtagagcauc atgatccugt

30

<210> 2
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 Nucleic Acid Ligand

<220>
 <221> modified_base
 <222> (4)..(29)

<223> C at positions 4, 8, 21 and 29 is
2'-fluoro-2'-deoxycytidine

<220>

<221> modified_base

<222> (6)..(30)

<223> U at positions 6, 20 and 30 is
2'-fluoro-2'-deoxyuridine

<220>

<221> modified_base

<222> (5)..(31)

<223> G at positions 5, 9, 17, and 31 is
2'-O-Methyl-2'-deoxyguanosine

<220>

<221> modified_base

<222> (22)

<223> A at position 22 is 2'-O-Methyl-2'-deoxyadenosine

<220>

<221> modified_base

<222> (1)..(30)

<223> The residues at positions 9 and 10 are connected
by a hexaethylene glycol phosphoramidite linker.
The residues at positions 21 and 22 are
connected by a hexaethylene glycol phosphoramidite
linker.

<220>

<221> modified_base

<222> (32)

<223> Nucleotide 32 is an inverted orientation T
(3'-3'-linked)

<400> 2

cagcguaacgc gtaccgatuc atgaagcugt

30